

| Replicate No. at Each Station | Batch No. | Phylum | Class | Order | Family | Scientific Name | Quantity | Weight (g) | Average Weight (g) |
|-------------------------------|-----------|------------|--------------|---------------------|----------------|-----------------------------------|----------|------------|--------------------|
| 1-A | 7565-1 | Annelida | Polychaeta | Phyllodocida | Glyceridae | <i>Glycera onomichiensis</i> | 1 | 0.4780 | 0.4780 |
| | | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 2 | 8.9542 | 4.4771 |
| 1-B | 7565-2 | Annelida | Polychaeta | Phyllodocida | Glyceridae | <i>Glycera onomichiensis</i> | 2 | 0.3762 | 0.1881 |
| | | Chordata | Osteichthyes | Perciformes | Taenioididae | <i>Trypauchen vagina</i> | 1 | 0.0349 | 0.0349 |
| | | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 7 | 13.5211 | 1.9316 |
| 1-C | 7565-3 | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 1 | 2.9423 | 2.9423 |
| 1-D | 7565-4 | Annelida | Polychaeta | Phyllodocida | Glyceridae | <i>Glycera onomichiensis</i> | 1 | 0.1860 | 0.1860 |
| | | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 2 | 6.9350 | 3.4675 |
| 1-E | 7565-5 | Annelida | Polychaeta | Eunicida | Eunicidae | <i>Eunice indica</i> | 3 | 0.0222 | 0.0074 |
| | | Arthropoda | Crustacea | Decapoda | Alpheidae | <i>Alpheus</i> sp. | 1 | 0.1994 | 0.1994 |
| | | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 9 | 19.6441 | 2.1827 |
| | | Mollusca | Gastropoda | Sorbeoconcha | Nassariidae | <i>Nassarius festivus</i> | 1 | 0.1991 | 0.1991 |
| 1-F | 7565-6 | Annelida | Polychaeta | Phyllodocida | Glyceridae | <i>Glycera onomichiensis</i> | 1 | 0.2261 | 0.2261 |
| | | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 30 | 90.7563 | 3.0252 |
| 1-G | 7565-7 | Annelida | Polychaeta | Eunicida | Eunicidae | <i>Eunice indica</i> | 3 | 0.1612 | 0.0537 |
| | | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 29 | 110.5249 | 3.8112 |
| | | Mollusca | Gastropoda | Sorbeoconcha | Nassariidae | <i>Nassarius dorsatus</i> | 1 | 1.4472 | 1.4472 |
| | | Mollusca | Gastropoda | Sorbeoconcha | Nassariidae | <i>Nassarius festivus</i> | 1 | 0.5878 | 0.5878 |
| 1-H | 7565-8 | Annelida | Polychaeta | Capitellida | Capitellidae | <i>Mediomastus californiensis</i> | 4 | 0.0392 | 0.0098 |
| | | Annelida | Polychaeta | Eunicida | Eunicidae | <i>Eunice indica</i> | 2 | 0.0599 | 0.0300 |
| | | Annelida | Polychaeta | Phyllodocida | Glyceridae | <i>Glycera onomichiensis</i> | 3 | 0.6970 | 0.2323 |
| | | Mollusca | Bivalvia | Eulamellibranchiata | Mesodesmatidae | <i>Caecella chinensis</i> | 1 | 6.9818 | 6.9818 |
| | | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 2 | 6.6248 | 3.3124 |

Sediment Size Distribution

| Parameters | | Station 1 |
|--|------------------------------------|-----------|
| Size Spectrum (Percentage of sand/silt/clay) | Size Spectrum (>2000 μ m), % | 0.7~1.9 |
| | Size Spectrum (63-2000 μ m), % | 15~26 |
| | Size Spectrum (<63 μ m), % | 72~84 |

Remarks:

1. Analytical Method - GEOSPEC 3:2001 Test 8.1