

| Replicate No. at Each Station | Batch No. | Phylum | Class | Order | Family | Scientific Name | Quantity | Weight (g) | Average Weight (g) |
|-------------------------------|-----------|----------|----------|---------------------|--------------|--------------------------------|----------|------------|--------------------|
| 2-A | 7565-9 | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 5 | 1.1650 | 0.2330 |
| 2-B | 7565-10 | none | | | | | | | |
| 2-C | 7565-11 | none | | | | | | | |
| 2-D | 7565-12 | none | | | | | | | |
| 2-E | 7565-13 | Mollusca | Bivalvia | Eulamellibranchiata | Psammobiidae | <i>Asaphis dichotoma</i> | 1 | 0.2486 | 0.2486 |
| | | Mollusca | Bivalvia | Veneroida | Veneridae | <i>Ruditapes philippinarum</i> | 11 | 2.2283 | 0.2026 |
| 2-F | 7565-14 | none | | | | | | | |
| 2-G | 7565-15 | none | | | | | | | |
| 2-H | 7565-16 | none | | | | | | | |

Sediment Size Distribution

| Parameters | | Station 2 |
|--|------------------------------------|-----------|
| Size Spectrum (Percentage of sand/silt/clay) | Size Spectrum (>2000 μ m), % | 0.5~1.1 |
| | Size Spectrum (63-2000 μ m), % | 8.6~13 |
| | Size Spectrum (<63 μ m), % | 86~91 |

Remarks:

1. Analytical Method - GEOSPEC 3:2001 Test 8.1