

Results of Benthic Survey of September 2011

Site	Phylum	Class	Order	Family	Scientific Name	Quantity	Weight (g)	Average weight (g)
3A	Annelida	Polychaeta	Terebellida	Pectinariidae	<i>Amphictene japonica</i>	2	0.0366	0.0183
3B	Annelida	Polychaeta	Phyllodocida	Nereididae	<i>Tylorrhynchus heterochaetus</i>	2	0.0230	0.0115
3C	None	--	--	--	--	--	--	--
3D	Annelida	Polychaeta	Aciculata	Onuphidae	<i>Diopatra</i> spp.	1	0.2067	0.2067
	Mollusca	Gastropoda	Neogastropoda	Nassariidae	<i>Nassarius siquijorensis</i>	2	0.4226	0.2113
3E	Annelida	Polychaeta	Orbiniida	Orbiniidae	<i>Scoloplos armiger</i>	2	0.0080	0.0040
3F	None	--	--	--	--	--	--	--
3G	Annelida	Polychaeta	Phyllodocida	Glyceridae	<i>Glycera onomichiensis</i>	1	0.1627	0.1627
3H	Annelida	Polychaeta	Scolecida	Paraonidae	<i>Paradoneis</i> spp.	2	0.0124	0.0062
	Annelida	Polychaeta	Aciculata	Nephtyidae	<i>Aglaophamus inermis</i>	2	0.0066	0.0033

Sediment Size Distribution

		Range
Parameter	Station 3	
Size Spectrum (Percentage of sand/silt/clay)	Size Spectrum (>2000µm), %	0.9
	Size Spectrum (63-2000µm), %	15
	Size Spectrum (<63µm), %	84-85