

Results of Benthic Survey of September 2011

Site	Phylum	Class	Order	Family	Scientific Name	Quantity	Weight (g)	Average weight (g)
SS3A	None	--	--	--	--	--	--	--
SS3B	None	--	--	--	--	--	--	--
SS3C	Annelida	Polychaeta	Aciculata	Nephtyidae	<i>Aglaophamus inermis</i>	27	0.0995	0.0037
	Annelida	Polychaeta	Orbiniida	Orbiniidae	<i>Naineris laevigata</i>	9	0.0438	0.0049
	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Ruditapes philippinarum</i>	2	0.0263	0.0132
SS3D	Annelida	Polychaeta	Aciculata	Nephtyidae	<i>Aglaophamus inermis</i>	19	0.1033	0.0054
	Annelida	Polychaeta	Terebellida	Pectinariidae	<i>Amphictene japonica</i>	2	0.0044	0.0022
SS3E	Annelida	Polychaeta	Aciculata	Phyllodocidae	<i>Eteone</i> spp.	7	0.0250	0.0036
	Annelida	Polychaeta	Scolecida	Capitellidae	<i>Mediomastus californiensis</i>	1	0.0024	0.0024
	Annelida	Polychaeta	Orbiniida	Orbiniidae	<i>Naineris laevigata</i>	5	0.0157	0.0031
SS3F	None	--	--	--	--	--	--	--
SS3G	Annelida	Polychaeta	Scolecida	Maldanidae	<i>Praxillela</i> spp.	1	0.0582	0.0582
	Annelida	Polychaeta	Scolecida	Capitellidae	<i>Mediomastus californiensis</i>	2	0.0181	0.0091
	Annelida	Polychaeta	Aciculata	Eunicidae	<i>Eunice indica</i>	1	0.0030	0.0030
	Annelida	Polychaeta	Aciculata	Nephtyidae	<i>Aglaophamus inermis</i>	14	0.0494	0.0035
SS3H	Annelida	Polychaeta	Scolecida	Capitellidae	<i>Mediomastus californiensis</i>	1	0.0013	0.0013
	Annelida	Polychaeta	Aciculata	Eunicidae	<i>Eunice indica</i>	1	0.0007	0.0007
	Mollusca	Gastropoda	Neogastropoda	Nassariidae	<i>Nassarius siquijorensis</i>	1	0.0681	0.0681

Sediment Size Distribution

Parameter	Range	
	Station SS3	
Size Spectrum (Percentage of sand/silt/clay)	Size Spectrum (>2000µm), %	0.7
	Size Spectrum (63-2000µm), %	14~16
	Size Spectrum (<63µm), %	84~85