

**Results of Benthic Survey of September 2011**

Site	Phylum	Class	Order	Family	Scientific Name	Quantity	Weight (g)	Average weight (g)
VS6A	Mollusca	Gastropoda	Neogastropoda	Nassariidae	<i>Nassarius festivus</i>	1	0.2098	0.2098
VS6B	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Ruditapes philippinarum</i>	2	0.7859	0.3930
VS6C	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Ruditapes philippinarum</i>	2	1.1470	0.5735
VS6D	Mollusca	Gastropoda	Neogastropoda	Nassariidae	<i>Nassarius festivus</i>	4	0.5991	0.1498
	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Ruditapes philippinarum</i>	7	4.6621	0.6660
VS6E	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Ruditapes philippinarum</i>	4	1.5478	0.3870
VS6F	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Ruditapes philippinarum</i>	5	1.6533	0.3307
VS6G	None	--	--	--	--	--	--	--
VS6H	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Ruditapes philippinarum</i>	3	2.4703	0.8234
	Annelida	Polychaeta	Phyllodocida	Glyceridae	<i>Glycera onomichiensis</i>	1	0.0059	0.0059
	Annelida	Polychaeta	Solecida	Opheliidae	<i>Ophelina grandis</i>	1	0.0042	0.0042
	Annelida	Polychaeta	Scolecida	Capitellidae	<i>Mediomastus californiensis</i>	4	0.0998	0.0250

**Sediment Size Distribution**

Parameter	Range	
	Station VS6	
Size Spectrum (Percentage of sand/silt/clay)	Size Spectrum (>2000µm), %	5.1~5.5
	Size Spectrum (63-2000µm), %	14~15
	Size Spectrum (<63µm), %	80